

# Septicaemia

Some bacteria can cause meningitis and septicaemia (blood poisoning). People with septicaemia may develop a rash of tiny red 'pin pricks' which can develop into purple bruising.

**THIS RASH DOES NOT FADE UNDER PRESSURE. DO THE GLASS TEST.**



Press the side of a clear glass firmly against the skin.

Spots/rash may fade at first.

Keep checking!

- Fever with spots/rash that do not fade under pressure is a **MEDICAL EMERGENCY**
- **DO NOT WAIT FOR A RASH.** If someone is ill and getting worse, get medical help immediately
- **ON DARK SKIN**, the spots/rash can be more difficult to see. Do not wait for a rash. Be aware of all the signs and symptoms.

# Getting medical help

If you are concerned it could be meningitis or septicaemia, you can:

Call NHS 111 or your GP

In an emergency you can:

- Dial 999 for an ambulance
- Go to your nearest accident and emergency department

Describe the symptoms and say that you think it could be meningitis or septicaemia.

Early diagnosis can be difficult. If you have had advice and are still concerned, get medical help again.



Download our free app  
Available on iPhone and Android

[www.meningitisapp.co.uk](http://www.meningitisapp.co.uk)

# Keep this card safe

## It could save a life

Common signs and symptoms for meningitis and septicaemia

**Meningitis  
NOW**

Meningitis Helpline  
**0808 80 10 388 (UK)**

[www.MeningitisNow.org](http://www.MeningitisNow.org)



**Supported by the Petra Stunt Foundation**

© Meningitis Now July 2017. Next review June 2018.  
References for content of this card are available on our website.  
Reg. Charity Number 803016 (England & Wales) SC037790 (Scotland)  
Company Registration Number 2469130.

## Babies and Toddlers



Fever, cold hands and feet



Refusing food and vomiting



Fretful, dislike being handled



Drowsy, floppy, unresponsive



Rapid breathing or grunting



Pale, blotchy skin  
Spots/rash  
**See Glass Test**



Unusual cry, moaning



Tense, bulging fontanelle (soft spot)



Stiff neck, dislike bright lights



Convulsions/seizures

## Be aware

Meningitis and septicaemia often happen together. Be aware of all the signs and symptoms.

Symptoms can appear in any order.

Some may not appear at all.

Early symptoms can include:

fever headache vomiting diarrhoea

muscle pain stomach cramps

fever with cold hands and feet

Someone with meningitis or septicaemia can get a lot worse very quickly. Keep checking them.

Trust your instincts -  
get medical help  
immediately



## Children and Adults



Fever, cold hands and feet



Vomiting



Drowsy, difficult to wake



Confusion and irritability



Severe muscle pain



Pale, blotchy skin  
Spots/rash  
**See Glass Test**



Severe headache



Stiff neck



Dislike bright lights



Convulsions/seizures